



LIFELINE  
SPECIALTY PHARMACY

PROVIDES:

Benefits  
investigation

Prior authorization  
and appeals  
support

Fulfillment with  
free shipping

# Send PANCREAZE prescriptions to LifeLine Specialty Pharmacy



USE THE FOLLOWING TO PRESCRIBE:



By Fax

**TO:** LifeLine Specialty  
Pharmacy

**NCPDP:** 2141240

**NPI:** 1568975464

**FAX:** 833-785-4461



e-Prescribe

**TO:** LifeLine Specialty  
Pharmacy

**NCPDP:** 2141240

**NPI:** 1568975464



Instructions

A patient's dose should be **individualized** based on age, weight, clinical symptoms, degree of steatorrhea, and fat content of their diet.

**Prescribe PANCREAZE for both meals and snacks** — usually, half of the prescribed dose for an individualized full meal should be given with each snack.

TOLL FREE

1-833  
454-3354

(4-LIFELINE)

**LOCATION:** 6304 Woodside Ct, Ste 100, Columbia, Maryland 21046

LifeLine Specialty Pharmacy will contact your patient directly. Please provide the LifeLine Specialty Pharmacy Patient Information Card located at [HCP.PANCREAZE.com/materials](http://HCP.PANCREAZE.com/materials).

**Questions about dosing?** Please consult guides and dosing calculator on HCP website.



LifeLine Specialty Pharmacy is NOT AVAILABLE to patients in Puerto Rico, Guam or the U.S. Virgin Islands.  
**See reverse for additional information and services.**



# PANCREAZE...

The symptoms of EPI<sup>SM</sup>



- Proven to improve EPI symptoms and fat and protein absorption.<sup>1,2</sup>
- Available in 6 strengths — including a 37,000 unit dose.
- A trusted enzyme replacement therapy for over 15 years.

### Important Safety Information

Fibrosing Colonopathy: Associated with high doses, usually over prolonged use and in pediatric patients with cystic fibrosis. Colonic stricture reported in pediatric patients less than 12 years of age with dosages exceeding 6,000 lipase units/kg/meal. Monitor during treatment for progression of preexisting disease. Do not exceed the recommended dosage, unless clinically indicated.

Hyperuricemia has been reported with high dosages; consider monitoring blood uric acid levels in patients with gout, renal impairment, or hyperuricemia.

Irritation of the oral mucosa may occur due to loss of protective enteric coating on the capsule contents.

The presence of porcine viruses that might infect humans cannot be definitely excluded.

Monitor patients with known reactions to proteins of porcine origin. If symptoms occur, initiate appropriate medical management; consider the risks and benefits of continued treatment.

### References:

1. PANCREAZE Full Prescribing Information. Campbell, CA: VIVUS LLC; 2024. 2. Trapnell BC, et al. Efficacy and safety of PANCREAZE<sup>®</sup> for treatment of exocrine pancreatic insufficiency due to cystic fibrosis. *J Cyst Fibros*. 2011;10(5):350-356.

Please read the accompanying **PANCREAZE Medication Guide** and **Full Prescribing Information**.

©2026 VIVUS LLC. All rights reserved.  
500392.04-USP 01/2026

For more information, visit: [HCP.PANCREAZE.com](https://HCP.PANCREAZE.com)

VIVUS